

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 June 2003 (19.06.2003)

PCT

(10) International Publication Number
WO 03/049724 A1

(51) International Patent Classification⁷: **A61K 31/00**,
31/138, A61P 25/28

(21) International Application Number: PCT/US02/36132

(22) International Filing Date:
27 November 2002 (27.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/339,174 11 December 2001 (11.12.2001) US

(71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BYMASTER, Franklin, Porter** [US/US]; 8545 North 650 East, Brownsburg, IN 46112 (US). **GEHLERT, Donald, Richard** [US/US]; 1440 Broadway, Indianapolis, IN 46202 (US). **MCKINZIE, David, Lee** [US/US]; 3693 South 500 East, Whitestown, IN 46075 (US). **YANG, Charles, Renkin** [CA/US]; 6215 North Washington Boulevard, Indianapolis, IN 46220 (US).

(74) Agents: **TITUS, Robert, D.** et al.; Eli Lilly and Company, P.O. Box 6288, Indianapolis, IN 46206-6288 (US).

(81) Designated States (national): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,

MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW. ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF NOREPINEPHRINE REUPTAKE INHIBITORS FOR THE TREATMENT OF COGNITIVE FAILURE

(57) Abstract: Selective norepinephrine reuptake inhibitors, particularly atomoxetine, reboxetine and 2-alkylthio substituted phenoxyphenyl propylamines, are used for the treatment of cognitive failure, including cognitive failure due to dementia, delirium and schizophrenia.

WO 03/049724 A1

-1-

USE OF NOREPINEPHRINE REUPTAKE INHIBITORS FOR THE TREATMENT OF COGNITIVE FAILURE

The invention belongs to the fields of pharmaceutical chemistry and central nervous system medicine, and provides a method of treatment for cognitive failure.

5 Cognitive failure is the dysfunction or loss of cognitive functions, the processes by which knowledge is acquired, retained, and used. An estimated 2% of all Americans have some form and degree of cognitive failure, with about 15% of those over the age of 65 affected. Cognitive failure is most commonly due to delirium or
10 dementia, but it may also occur in association with a number of other disorders.

A delerium is characterized by a disturbance of consciousness and a change in cognition that develop over a short period of time. Dementia is a chronic deterioration
15 of intellectual function and other cognitive skills severe enough to interfere with the ability to perform activities of daily living. Although dementia may occur at any age, it primarily affects the elderly, presenting in more than 15% of persons over 65 years of age and in as many as 40% of
20 persons over 80 years old. Dementia accounts for more than half of nursing home admissions. Dementia due to Alzheimer's disease affects four million Americans, at an annual cost of about \$90 billion, including medical and
25 nursing home care, social services, lost productivity, and early death. Alzheimer's disease accounts for more than 65% of the dementias in the elderly.

Non-Alzheimer's dementias include Lewy body dementia, vascular dementia, and Binswanger's dementia
30 (subcortical arteriosclerotic encephalopathy). Dementia may also appear in patients with Parkinson's disease, progressive supranuclear palsy, Huntington's disease (chorea), Pick's disease, frontal lobe dementia syndromes, dementia pugilistica, normal-pressure hydrocephalus,

-2-

subdural hematoma, Creutzfeldt-Jakob disease, Gerstmann-Sträussler-Scheinker disease, general paresis, AIDS, and schizophrenia.

Current treatments for cognitive failure include compounds that enhance cholinergic transmission such as donepezil, rivastigmine, and tacrine. The use of these agents is limited by side effects including changes in vision or balance, diarrhea, dizziness, fainting spells or falls, increase in frequency of passing urine or incontinence, nervousness, agitation, increased confusion, skin rash or hives, slow heartbeat or palpitations, stomach pain, sweating, uncontrollable movements, unusual bleeding or bruising, red or purple spots on the skin, vomiting, and weight loss. Another therapy is the administration of the ergot hydergine. Hydergine therapy may require six months to determine whether the drug has been effective, and side effects include nausea.

Additional therapies are needed for the treatment of cognitive failure that are more efficacious and better tolerated than treatments that are currently available.

The present invention provides a method for the treatment of cognitive failure that comprises administering to a mammal in need of such treatment an effective amount of a selective norepinephrine reuptake inhibitor.

The present invention provides a method for the treatment of cognitive failure that relies on a novel mechanism of action. This method comprises treating a mammal suffering from cognitive failure with a compound that is a selective norepinephrine reuptake inhibitor. This mechanism is operative in mammals and the preferred mammal is a human.

The present invention also provides the use of a selective norepinephrine reuptake inhibitor for the

-3-

preparation of a medicament useful for the treatment or prevention of cognitive failure.

The present invention provides a method for the treatment of cognitive failure. The term "cognitive failure", as used herein, refers to the spectrum of cognitive dysfunctions ranging from mild cognitive impairment to deterioration of intellectual function and other cognitive skills severe enough to interfere with the ability to perform activities of daily living.


Many compounds, including those discussed at length below, are selective norepinephrine reuptake inhibitors, and no doubt many more will be identified in the future. In the practice of the present invention, it is intended to include reuptake inhibitors which show 50% effective concentrations of about 1000 nM or less, in the protocol described by Wong et al., *Drug Development Research*, 6, 397 (1985). The norepinephrine reuptake inhibitors useful for the method of the present invention are characterized in being selective for the inhibition of neurotransmitter reuptake relative to their ability to act as direct agonists or antagonists at other receptors. It is preferred that the compounds useful for the method of the present invention are selective for the inhibition of norepinephrine reuptake relative to direct agonist or antagonist activity at other receptors by a factor of at least ten. Preferably, compounds useful for the method of the present invention are selective for the inhibition of norepinephrine reuptake relative to direct agonist or antagonist activity at other receptors by a factor of at least one hundred. Norepinephrine reuptake inhibitors useful for the method of the present invention include, but are not limited to:

Atomoxetine (formerly known as tomoxetine), (R)-

-4-

(-)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine, is usually administered as the hydrochloride salt. Atomoxetine was first disclosed in U.S. Patent #4,314,081. The word "atomoxetine" will be used here to refer to any acid addition salt or the free base of the molecule. See, for example, Gehlert, et al., *Neuroscience Letters*, 157, 203-206 (1993), for a discussion of atomoxetine's activity as a norepinephrine reuptake inhibitor;

The compounds of formula I:



I

wherein X is C₁-C₄ alkylthio, and Y is C₁-C₂ alkyl or a pharmaceutically acceptable salt thereof. The compounds of formula I were described in U.S. Patent 5,281,624, of Gehlert, Robertson, and Wong, and in Gehlert, et al., *Life Sciences*, 55(22), 1915-1920, (1995). The compounds are there taught to be inhibitors of norepinephrine reuptake in the brain. It is also explained that the compounds exist as stereoisomers, and that they accordingly include not only the racemates, but also the isolated individual isomers as well as mixtures of the individual isomers. For example, the compounds of formula I include the following exemplary species:

N-ethyl-3-phenyl-3-(2-methylthiophenoxy)propylamine benzoate;

(R)-N-methyl-3-phenyl-3-(2-propylthiophenoxy)-propylamine hydrochloride;

-5-

(S)-N-ethyl-3-phenyl-3-(2-butylthiophenoxy)propyl-amine;

N-methyl-3-phenyl-3-(2-ethylthiophenoxy)propyl-amine malonate;

5 (S)-N-methyl-3-phenyl-3-(2-tert-butylthiophenoxy)-propylamine naphthalene-2-sulfonate;

(R)-N-methyl-3-(2-methylthiophenoxy)-3-phenyl-propylamine; and

10 Reboxetine (EdronaxTM), 2-[α -(2-ethoxy)phenoxy-benzyl]morpholine, is usually administered as the racemate. It was first taught by U.S. Patent 4,229,449, which describes its utility for the treatment of depression. Reboxetine is a selective norepinephrine reuptake inhibitor. The term "reboxetine" will be used here to refer to any acid
15 addition salt or the free base of the molecule existing as the racemate or either enantiomer.

While all compounds exhibiting norepinephrine reuptake inhibition are useful for the methods of the present invention, certain are preferred. It is preferred
20 that the norepinephrine reuptake inhibitor is selective for the reuptake of norepinephrine over the reuptake of other neurotransmitters. It is also preferred that the norepinephrine reuptake inhibitor does not exhibit significant direct agonist or antagonist activity at other
25 receptors. It is especially preferred that the norepinephrine reuptake inhibitor be selected from atomoxetine, reboxetine, or (R)-N-methyl-3-(2-methylthio-phenoxy)-3-phenylpropylamine. The use of atomoxetine hydrochloride for the methods of the present invention is
30 the most preferred embodiment of the present invention.

A further embodiment of this invention comprises the administration of a composition that exhibits selective norepinephrine reuptake inhibitor activity. The composition may be composed of one or more agents that, individually or

-6-

together, are selective inhibitors of norepinephrine reuptake.

The dosages of the drugs used in the present invention must, in the final analysis, be set by the physician in charge of the case using knowledge of the drugs, the properties of the drugs in combination as determined in clinical trials, and the characteristics of the patient including diseases other than that for which the physician is treating the patient. General outlines of the dosages, and some preferred dosages, can and will be provided here.

Atomoxetine: In adults and older adolescents: from about 5 mg/day to about 200 mg/day; preferably in the range from about 60 to about 150 mg/day; more preferably from about 60 to about 130 mg/day; and still more preferably from about 50 to about 120 mg/day; in children and younger adolescents: from about 0.2 to about 3.0 mg/kg/day; preferably in the range from about 0.5 to about 1.8 mg/kg/day;

Compounds of formula I: from about 0.01 mg/kg to about 20 mg/kg; preferred daily doses will be from about 0.05 mg/kg to 10 mg/kg; ideally from about 0.1 mg/kg to about 5 mg/kg;

Reboxetine: from about 1 to about 30 mg, once to four times/day; preferred, from about 5 to about 30 mg once/day.

Cognitive failure presents in patients suffering from a number of other disorders. The present invention includes the use of a norepinephrine reuptake inhibitor to treat cognitive failure presenting alone or where cognitive failure is associated with another disorder. Schizophrenic patients, for example, commonly exhibit symptoms that include cognitive failure. An embodiment of the present invention, therefore, is the use of a norepinephrine

-7-

reuptake inhibitor to treat cognitive failure associated with schizophrenia. Patients suffering from schizophrenia also frequently exhibit negative symptoms such as flat affect, asociality, anergia, avolition, and anhedonia. A further embodiment of the present invention is the use of a norepinephrine reuptake inhibitor to treat the negative symptoms of schizophrenia.

The invention further provides a method for treating a patient suffering from or susceptible to psychosis, comprising administering to said patient an effective amount of a first component which is an antipsychotic, in combination with an effective amount of a second component which is a norepinephrine reuptake inhibitor. The invention also provides a pharmaceutical composition that comprises a first component that is an antipsychotic, and a second component that is a norepinephrine reuptake inhibitor.

In the general expressions of this aspect of the present invention, the first component is a compound that acts as an antipsychotic. The antipsychotic may be either a typical antipsychotic, such as haloperidol, or an atypical antipsychotic. The essential feature of an atypical antipsychotic is less acute extrapyramidal symptoms, especially dystonias, associated with therapy as compared to a typical antipsychotic such as haloperidol. Clozapine, the prototypical atypical antipsychotic, differs from the typical antipsychotics with the following characteristics:

- (1) greater efficacy in the treatment of overall psychopathology in patients with schizophrenia nonresponsive to typical antipsychotics;
- (2) greater efficacy in the treatment of negative symptoms of schizophrenia; and
- (3) less frequent and quantitatively smaller increases in serum prolactin concentrations associated with therapy (Beasley, et al., Neuropsychopharmacology, 14(2), 111-123, (1996)).

-8-

Although both typical and atypical antipsychotics are useful for these methods and formulations of the present invention, it is preferred that the first component compound is an atypical antipsychotic.

5 Typical antipsychotics include, but are not limited to:

 Chlorpromazine, 2-chloro-10-(3-dimethylaminoprop-
yl)phenothiazine, is described in U.S. Patent 2,645,640.
Its pharmacology has been reviewed (Crismon, Psychopharma-
10 col. Bul., 4, 151 (October 1967));

 Droperidol, 1-(1-[3-(p-fluorobenzoyl)propyl]-
1,2,3,6-tetrahydro-4-pyridyl)-2-benzimidazolinone, is
described in U.S. Patent 3,141,823;

 Haloperidol, 4-[4-(4-chlorophenyl)-4-hydroxy-1-
15 piperidinyl]-1-(4-fluorophenyl)-1-butanone, is described in
U.S. Patent 3,438,991. Its therapeutic efficacy in
psychosis has been reported (Beresford and Ward, Drugs, 33,
31-49 (1987));

 Thioridazine, 1-hydroxy-10-[2-(1-methyl-2-
20 pyridinyl)ethyl]-2-(methylthio)phenothiazine hydrochloride,
was described by Bourquin, et al. (Helv. Chim. Acta, 41, 1072
(1958)). Its use as an antipsychotic has been reported
(Axelsson, et al., Curr. Ther. Res., 21, 587 (1977)); and

 Trifluoperazine, 10-[3-(4-methyl-1-piperazinyl)-
25 propyl]-2-trifluoromethylphenthiazine hydrochloride, is
described in U.S. Patent 2,921,069.

 Atypical antipsychotics include, but are not
limited to: Olanzapine, 2-methyl-4-(4-methyl-1-
piperazinyl)-10H-thieno[2,3-b][1,5]benzodiazepine, is a
30 known compound and is described in U.S. Patent No. 5,229,382
as being useful for the treatment of schizophrenia,
schizophreniform disorder, acute mania, mild anxiety states,
and psychosis;

-9-

Clozapine, 8-chloro-11-(4-methyl-1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine, is described in U.S. Patent No. 3,539,573. Clinical efficacy in the treatment of schizophrenia is described (Hanes, et al., Psychopharmacol. Bull., 24, 62 (1988));

Risperidone, 3-[2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidino]ethyl]-2-methyl-6,7,8,9-tetrahydro-4H-pyrido[1,2-a]pyrimidin-4-one, and its use in the treatment of psychotic diseases are described in U.S. Patent No. 4,804,663;

Sertindole, 1-[2-[4-[5-chloro-1-(4-fluorophenyl)-1H-indol-3-yl]-1-piperidinyl]ethyl]imidazolidin-2-one, is described in U.S. Patent No. 4,710,500. Its use in the treatment of schizophrenia is described in U.S. Patent Nos. 5,112,838 and 5,238,945;

Quetiapine, 5-[2-(4-dibenzo[b,f][1,4]thiazepin-11-yl-1-piperazinyl)ethoxy]ethanol, and its activity in assays which demonstrate utility in the treatment of schizophrenia are described in U.S. Patent No. 4,879,288. Quetiapine is typically administered as its (E)-2-butenedioate (2:1) salt; and

Ziprasidone, 5-[2-[4-(1,2-benzisothiazol-3-yl)-1-piperazinyl]ethyl]-6-chloro-1,3-dihydro-2H-indol-2-one, is typically administered as the hydrochloride monohydrate. The compound is described in U.S. Patent Nos. 4,831,031 and 5,312,925. Its activity in assays which demonstrate utility in the treatment of schizophrenia are described in U.S. Patent No. 4,831,031.

Similarly, when this aspect of the invention is regarded in its broadest sense, the second component compound is a compound that functions as a norepinephrine reuptake inhibitor as described above.

It will be understood that while the use of a single antipsychotic as a first component compound is

-10-

preferred, combinations of two or more antipsychotics may be used as a first component if necessary or desired.

Similarly, while the use of a single norepinephrine reuptake inhibitor as a second component compound is preferred,

5 combinations of two or more norepinephrine reuptake inhibitors may be used as a second component if necessary or desired.

While all combinations of first and second component compounds are useful and valuable, certain
10 combinations are particularly valued and are preferred, as follows:

olanzapine/atomoxetine

olanzapine/reboxetine

olanzapine/(R)-N-methyl-3-(2-methylthiophenoxy)-3-
15 phenylpropylamine

clozapine/atomoxetine

risperidone/atomoxetine

sertindole/atomoxetine

quetiapine/atomoxetine

20 ziprasidone/atomoxetine

In general, combinations and methods of treatment using olanzapine as the first component are preferred.

Furthermore, combinations and methods of treatment using atomoxetine as the second component are preferred.

25 Especially preferred are combinations and methods of treatment using olanzapine as the first component and atomoxetine as the second component. It is especially preferred that when the first component is olanzapine, it will be the Form II olanzapine as described in U.S. Patent
30 5,736,541.

It is further preferred that the Form II olanzapine polymorph will be administered as the substantially pure Form II olanzapine polymorph. As used herein "substantially pure" refers to Form II associated

-11-

with less than about 5% Form I, preferably less than about 2% Form I, and more preferably less than about 1% Form I. Further, "substantially pure" Form II will contain less than about 0.5% related substances, wherein "related substances" refers to undesired chemical impurities or residual solvent or water. In particular, "substantially pure" Form II should contain less than about 0.05% content of acetonitrile, more preferably, less than about 0.005% content of acetonitrile. Additionally, the polymorph of the invention should contain less than 0.5% of associated water.

Although Form II olanzapine is preferred it will be understood that as used herein, the term "olanzapine" embraces all solvate and polymorphic forms unless specifically indicated.

It will be understood by the skilled reader that most or all of the compounds used in the present invention are capable of forming salts, and that the salt forms of pharmaceuticals are commonly used, often because they are more readily crystallized and purified than are the free bases. In all cases, the use of the pharmaceuticals described above as salts is contemplated in the description herein, and often is preferred, and the pharmaceutically acceptable salts of all of the compounds are included in the names of them. Especially preferred pharmaceutically acceptable salts are those formed with hydrochloric acid.

The dosages of the first component drugs used in this aspect of the present invention must, in the final analysis, be set by the physician in charge of the case, using knowledge of the drugs, the properties of the drugs in combination as determined in clinical trials, and the characteristics of the patient, including diseases other than that for which the physician is treating the patient. General outlines of the dosages, and some preferred dosages, can and will be provided here. Dosage guidelines for some

-12-

of the drugs will first be given separately; in order to create a guideline for any desired combination, one would choose the guidelines for each of the component drugs.

Chlorpromazine: from about 25-75 mg daily to about 75-150 mg daily;

Droperidol: about 5 mg by injection;

Haloperidol: from about 1-15 mg/day to about 100 mg/day administered orally or by injection;

Thioridazine: about 75-150 mg daily;

Trifluoperazine: from about 4-10 mg/day to about 15-20 mg/day;

Olanzapine: from about 0.25 to 50 mg, once/day; preferred, from 1 to 30 mg, once/day; and most preferably 1 to 25 mg once/day;

Clozapine: from about 12.5 to 900 mg daily; preferred, from about 150 to 450 mg daily;

Risperidone: from about 0.25 to 16 mg daily; preferred from about 2-8 mg daily;

Sertindole: from about .0001 to 1.0 mg/kg daily;

Quetiapine: from about 1.0 to 40 mg/kg given once daily or in divided doses;

Ziprasidone: from about 5 to 500 mg daily; preferred from about 50 to 100 mg daily.

In more general terms, one would create a combination of the present invention by choosing a dosage of first and second component compounds according to the spirit of the above guideline.

The adjunctive therapy aspect of the present invention is carried out by administering a first component together with the second component in any manner that provides effective levels of the compounds in the body at the same time. Oral administration of the adjunctive combination is preferred. Both components may be administered together, in a single dosage form, or may be

-13-

administered separately. However, oral administration is not the only route or even the only preferred route. For example, transdermal administration may be very desirable for patients who are forgetful or petulant about taking oral medicine. Administration by the percutaneous, intravenous, intramuscular, intranasal or intrarectal route may be prudent in particular circumstances. The route of administration may be varied in any way, limited by the physical properties of the drugs, the convenience of the patient and the caregiver, and other relevant circumstances (Remington's Pharmaceutical Sciences, 18th Edition, Mack Publishing Co. (1990)).

The adjunctive combination may be administered as a single pharmaceutical composition, and so pharmaceutical compositions incorporating both compounds are important embodiments of the present invention. Such compositions may take any physical form that is pharmaceutically acceptable, but orally usable pharmaceutical compositions are particularly preferred. Such adjunctive pharmaceutical compositions contain an effective amount of each of the compounds, which effective amount is related to the daily dose of the compounds to be administered. Each adjunctive dosage unit may contain the daily doses of all compounds, or may contain a fraction of the daily doses, such as one-third of the doses. Alternatively, each dosage unit may contain the entire dose of one of the compounds, and a fraction of the dose of the other compounds. In such case, the patient would daily take one of the combination dosage units, and one or more units containing only the other compounds. The amounts of each drug to be contained in each dosage unit depends on the identity of the drugs chosen for the therapy, and other factors such as the indication for which the adjunctive therapy is being given.

-14-

The pharmaceutical compositions are prepared in a manner well known in the pharmaceutical art. The carrier or excipient may be a solid, semi-solid, or liquid material that can serve as a vehicle or medium for the active ingredient. Suitable carriers or excipients are well known in the art. The pharmaceutical composition may be adapted for oral, inhalation, parenteral, or topical use and may be administered to the patient in the form of tablets, capsules, aerosols, inhalants, suppositories, solutions, suspensions, or the like.

The compounds useful for the method of the present invention may be administered orally, for example, with an inert diluent or capsules or compressed into tablets. For the purpose of oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like. These preparations should contain at least 4% of the compound of the present invention, the active ingredient, but may be varied depending upon the particular form and may conveniently be between 4% to about 70% of the weight of the unit. The amount of the compound present in compositions is such that a suitable dosage will be obtained. Preferred compositions and preparations useful for the methods of the present invention may be determined by a person skilled in the art.

The tablets, pills, capsules, troches, and the like may also contain one or more of the following adjuvants: binders such as microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch or lactose, disintegrating agents such as alginic acid, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; and sweetening agents such as sucrose or saccharin may be added

-15-

or a flavoring agent such as peppermint, methyl salicylate or orange flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or a fatty oil. Other dosage unit forms may contain other various materials that modify the physical form of the dosage unit, for example, as coatings. Thus, tablets or pills may be coated with sugar, shellac, or other coating agents. A syrup may contain, in addition to the present compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

A formulation useful for the administration of R-(-)-N-methyl 3-((2-methylphenyl)oxy)-3-phenyl-1-aminopropane hydrochloride (atomoxetine) comprises a dry mixture of R-(-)-N-methyl 3-((2-methylphenyl)oxy)-3-phenyl-1-aminopropane hydrochloride with a diluent and lubricant. A starch, such as pregelatinized corn starch, is a suitable diluent and a silicone oil, such as dimethicone, a suitable lubricant for use in hard gelatin capsules. Suitable formulations are prepared containing about 0.4 to 26% R-(-)-N-methyl 3-((2-methylphen-yl)oxy)-3-phenyl-1-aminopropane hydrochloride, about 73 to 99% starch, and about 0.2 to 1.0% silicone oil. The following tables illustrate particularly preferred atomoxetine formulations:

-16-

Ingredient (%)	2.5 mg	5 mg	10 mg	18 mg	20 mg	25 mg	40 mg	60 mg
R-(-)-N-methyl 3- ((2-meth- ylphenyl)oxy)-3- phenyl-1- aminopropane hydrochloride	1.24	2.48	4.97	8.94	9.93	12.4 2	19.8 7	22.1 2
Dimethicone	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Pregelatinized Starch	98.2 6	97.0 2	94.5 3	90.5 6	89.5 7	87.0 8	79.6 3	77.3 8

Ingredient (mg/capsule)	2.5 mg	5 mg	10 mg	18 mg	20 mg	25 mg	40 mg	60 mg
R-(-)-N-methyl 3- ((2-meth- ylphenyl)oxy)-3- phenyl-1- aminopropane hydrochloride	2.86	5.71	11.4 3	20.5 7	22.8 5	28.5 7	45.7 1	68.5 6
Dimethicone	1.15	1.15	1.15	1.15	1.15	1.15	1.15	1.55
Pregelatinized Starch	225. 99	223. 14	217. 42	208. 28	206. 00	200. 28	183. 14	239. 89
Capsule Fill Weight (mg)	230	230	230	230	230	230	230	310
Capsule Size	3	3	3	3	3	3	3	2

5

For the purpose of parenteral therapeutic administration, the compounds of the present invention may be incorporated into a solution or suspension. These preparations typically contain at least 0.1% of a compound of the invention, but may be varied to be between 0.1 and

-17-

about 90% of the weight thereof. The amount of the compound of formula I present in such compositions is such that a suitable dosage will be obtained. The solutions or suspensions may also include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylene diaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Preferred compositions and preparations may be determined by one skilled in the art.

Inhibition or norepinephrine reuptake

The ability of compounds to inhibit the reuptake of norepinephrine may be measured by the general procedure of Wong, et al., *supra*.

Male Sprague-Dawley rats weighing 150-250 gm are decapitated and brains are immediately removed. Cerebral cortices are homogenized in 9 volumes of a medium containing 0.32 M sucrose and 10 mM glucose. Crude synaptosomal preparations are isolated after differential centrifugation at 1000 x g for 10 minutes and 17,000 x g for 28 minutes. The final pellets are suspended in the same medium and kept in ice until use within the same day.

Synaptosomal uptake of ^3H -norepinephrine is determined as follows. Cortical synaptosomes (equivalent to 1 mg of protein) are incubated at 37°C for 5 minutes in 1 mL Krebs-bicarbonate medium containing also 10 mM glucose, 0.1 mM iproniazide, 1 mM ascorbic acid, 0.17 mM EDTA and 50 nM

-18-

³H-norepinephrine. The reaction mixture is immediately diluted with 2 mL of ice-chilled Krebs-bicarbonate buffer and filtered under vacuum with a cell harvester (Brandel, Gaithersburg, MD). Filters are rinsed twice with approximately 5 mL of ice-chilled 0.9% saline and the uptake of ³H-norepinephrine assessed by liquid scintillation counting. Accumulation of ³H-norepinephrine at 4°C is considered to be background and is subtracted from all measurements. The concentration of the test compound required to inhibit 50% of the ³H-norepinephrine accumulation (IC₅₀ values) are determined by linear regression analysis.

The present invention provides a method for the treatment of cognitive failure. Cognitive failure may present in patients suffering from a number of disorders. The methods of the present invention are useful for the treatment of cognitive failure associated with disorders classified in the Diagnostic and Statistical Manual of Mental Disorders, 4th Version, published by the American Psychiatric Association (DSM-IV). The DSM code numbers are supplied below for the convenience of the reader.

	Delirium Due to a General Medical Condition	293.0
	Delirium Not Otherwise Specified	780.09
25	Dementia of the Alzheimer's Type	
	Early Onset with Delirium	290.11
	Early Onset with Delusions	290.12
	Early Onset Uncomplicated	290.10
	Late Onset with Delirium	290.3
30	Late Onset with Delusions	290.20
	Late Onset Uncomplicated	290.0
	Vascular Dementia	
	With Delirium	290.41
	With Delusions	290.42

-19-

	Uncomplicated	290.40
	Dementia Due to HIV Disease	294.1
	Dementia Due to Head Trauma	294.1
	Dementia Due to Parkinson's Disease	294.1
5	Dementia Due to Huntington's Disease	294.1
	Dementia Due to Pick's Disease	290.10
	Dementia Due to Creutzfeldt-Jakob Disease	290.10
	Dementia Due to Other General Medical Conditions	294.1
	Dementia Not Otherwise Specified	294.8
10	Amnestic Disorder Due to a General Medical Condition	294.0
	Amnestic Disorder Not Otherwise Specified	294.8
	Cognitive Disorder Not Otherwise Specified	294.9
	Paranoid Type Schizophrenia	295.30
15	Disorganized Type Schizophrenia	295.10
	Catatonic Type Schizophrenia	295.20
	Undifferentiated Type Schizophrenia	295.90
	Residual Type Schizophrenia	295.60
	Schizophreniform Disorder	295.40
20	Schizoaffective Disorder	295.70

The skilled artisan will appreciate that the disorders listed above are illustrative of those indications where cognitive failure may appear, and is not intended to limit the scope of the present invention in any way.

Psychotic conditions to be treated by the adjunctive therapy aspect of the present invention include schizophrenia, schizophreniform diseases, acute mania, and schizoaffective disorders. The titles given these conditions represent multiple disease states. The following list illustrates a number of these disease states, many of which are classified in the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, published by the American Psychiatric Association (DSM). The DSM code

-20-

numbers for these disease states are supplied below, when available, for the convenience of the reader.

Paranoid Type Schizophrenia	295.30
Disorganized Type Schizophrenia	295.10
Catatonic Type Schizophrenia	295.20
Undifferentiated Type Schizophrenia	295.90
Residual Type Schizophrenia	295.60
Schizophreniform Disorder	295.40
Schizoaffective Disorder	295.70

The present invention is also useful for the treatment of cognitive failure related to the onset of menopause.

The method of the present invention is effective in the treatment of patients who are children, adolescents or adults, and there is no significant difference in the symptoms or the details of the manner of treatment among patients of different ages. In general terms, however, for purposes of the present invention, a child is considered to be a patient below the age of puberty, an adolescent is considered to be a patient from the age of puberty up to about 18 years of age, and an adult is considered to be a patient of 18 years or older.

-21-

EXAMPLE 1

The immediate-early gene, c-fos, and its protein products have been increasingly utilized as markers for neuronal activation (Dragunow and Faull, J. Neurosci. Methods, **29**, 261-265 (1989); Morgan and Curran, Prog. In Brain Res., **86**, 287-294 (1990); Robertson, et al., J. Pharmacol. Exp. Ther., **271**, 1058-1066 (1994)). C-fos activation is measured as illustrated below for atomoxetine.

Two hours after administration of atomoxetine (3 mg/kg, i.p.), the rats were deeply anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and transcardially perfused with 100 ml of phosphate buffered saline (PBS) followed by 100 ml of 4% paraformaldehyde in PBS. The brain was rapidly removed, postfixed for 90 min in 4% paraformaldehyde and then was transferred to 30% sucrose at 4°C until saturated. After quick freezing, serial 30 µm sections were cut and placed in PBS until processed for immunohistochemistry. In brief, sections were incubated in PBS containing blocking serum and 0.5% Triton-X 100 for 1 hour. Sections were then incubated with anti-Fos antibody (Santa Cruz Biotechnology, Inc.) at 4°C overnight. Visualization of the Fos-like immunoreactivity was performed with a Vectastain ABC Elite Kit (Vector Labs, Burlingame, CA) using the standard protocol supplied with the kit. Nickel-intensified diaminobenzidine (DAB) was used as the chromagen to yield a gray-black precipitation product. Following visualization of the Fos immunoreactivity, the sections were mounted on gelatin-coated glass slides and allowed to dry. The sections were then dehydrated and cover slipped. Fos expressing cells were quantitated using the MCID M2 imaging system (Imaging Research, St. Catherines, Ontario).

Surprisingly, atomoxetine increased the expression of c-fos only in the cortical areas as demonstrated by the data in the following table.

-22-

Treatment	Prefrontal Cortex**	Nucleus Accumbens**	Striatum**
Vehicle	80 _± 28	129 _± 33	118 _± 26
Atomoxetine	296 _± 26*	152 _± 44	102 _± 35

*= $p < 0.001$ ** Fos positive cells/mm²


5.

-23-

We claim:

1. Use of a selective norepinephrine reuptake inhibitor for the manufacture of a medicament for the treatment of cognitive failure.

2. Use according to Claim 1 wherein the selective norepinephrine reuptake inhibitor is selected from the group consisting of atomoxetine, reboxetine, and a compound of formula I:



I

wherein X is C₁-C₄ alkylthio, and Y is C₁-C₂ alkyl or a pharmaceutically acceptable salt thereof.

3. Use according to Claim 2 wherein the selective norepinephrine reuptake inhibitor is atomoxetine.

4. Use according to Claim 2 wherein the selective norepinephrine reuptake inhibitor is atomoxetine hydrochloride.

5. Use according to any of Claims 1-4 wherein cognitive failure due to a dementia is treated.

6. Use according to any of Claims 1-4 wherein cognitive disorder due to a delirium is treated.

7. Use according to any of Claims 1-4 wherein cognitive failure due to schizophrenia is treated.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 02/36132

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/00 A61K31/138 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, MEDLINE, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 01973 A (MARSHALL ROBERT CLYDE ;UPJOHN CO (US); WONG ERIK H F (US); BIRGERS) 11 January 2001 (2001-01-11) page 1 -page 3 page 9 page 12	1,2,5-7
X,P	FERGUSON J M ET AL: "Effects of reboxetine on Hamilton Depression Rating Scale factors from randomized, placebo-controlled trials in major depression." INTERNATIONAL CLINICAL PSYCHOPHARMACOLOGY, vol. 17, no. 2, March 2002 (2002-03), pages 45-51, XP009008693 March, 2002 ISSN: 0268-1315 the whole document	1,2,5-7

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

31 March 2003

Date of mailing of the international search report

16/04/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Domingues, H

INTERNATIONAL SEARCH REPORT

Intern: I Application No
PCT/US 02/36132

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 721 777 A (LILLY CO ELI) 17 July 1996 (1996-07-17) the whole document ----	1-4
X,P	WO 02 053140 A (PHARMACIA AB ;SVENSSON TORGNY (SE); WONG ERIK HO FONG (US); UPJOHN) 11 July 2002 (2002-07-11) page 2 page 10 -page 15 ----	1-7
Y	GEHLERT D R ET AL: "(R)-THIONISOXETINE, A POTENT AND SELECTIVE INHIBITOR OF CENTRAL AND PERIPHERAL NOREPHINEPHRINE UPTAKE" LIFE SCIENCES, PERGAMON PRESS, OXFORD, GB, vol. 56, no. 22, 1995, pages 1915-1920, XP000602398 ISSN: 0024-3205 the whole document ----	1,2,5-7
X,P	WO 02 053104 A (EPSTEIN MEL H ;SENTION INC (US); WIIG KJESTEN A (US)) 11 July 2002 (2002-07-11) the whole document ----	1,2,5-7
Y	WO 01 27068 A (HOWARD HARRY RALPH JR ;PFIZER PROD INC (US); ADAM MAVIS DIANE (US)) 19 April 2001 (2001-04-19) the whole document ----	1,2,5-7
A	EP 0 534 756 A (LILLY CO ELI) 31 March 1993 (1993-03-31) the whole document -----	1-7

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/36132

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1 and 5-7
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 02 86132

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1 and 5-7

Present claims 1 and 5-7 relate to an extremely large number of possible compounds ("norepinephrine reuptake inhibitors") defined by reference to a desirable characteristic or property - the ability to inhibit the reuptake of norepinephrine, while support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds defined in claims 2-4 and for "norepinephrine reuptake inhibitor" in general. Attention is drawn to the fact that certain compounds might be already known for the treatment of cognitive disorders but might not have been identified as norepinephrine reuptake inhibitors.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 02 86132

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1, 2 (partially) and 3-7

1) Claims 1, 2 (partially) and 3-7 concern the use of the norepinephrine reuptake inhibitor atomoxetine for the treatment of cognitive failure due to dementia, delirium or schizophrenia.

2. Claims: 1, 2 (partially) and 5-7

2) Claims 1, 2 (partially) and 5-7 concern the use of the norepinephrine reuptake inhibitor reboxetine for the treatment of cognitive failure due to dementia, delirium or schizophrenia.

3. Claims: 1, 2 (partially) and 5-7

3) Claims 1, 2 (partially) and 5-7 concern the use of norepinephrine reuptake inhibitors of formula I for the treatment of cognitive failure due to dementia, delirium or schizophrenia.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/36132

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0101973	A	11-01-2001	AU 5633700 A	22-01-2001
			BR 0012136 A	11-06-2002
			CN 1379672 T	13-11-2002
			CZ 20014625 A3	14-08-2002
			EP 1196172 A2	17-04-2002
			HU 0201623 A2	28-09-2002
			JP 2003503450 T	28-01-2003
			NO 20016406 A	19-02-2002
			SK 19382001 A3	02-07-2002
			WO 0101973 A2	11-01-2001
			US 2002061910 A1	23-05-2002
			US 2002086864 A1	04-07-2002
			US 2002107249 A1	08-08-2002
			US 2002128173 A1	12-09-2002
			US 2003040464 A1	27-02-2003
			US 6465458 B1	15-10-2002
EP 0721777	A	17-07-1996	US 5658590 A	19-08-1997
			AT 222757 T	15-09-2002
			AU 688665 B2	12-03-1998
			AU 4693896 A	31-07-1996
			BR 9606903 A	21-10-1997
			CA 2209735 A1	18-07-1996
			CN 1168095 A	17-12-1997
			CZ 9702145 A3	14-01-1998
			DE 69623141 D1	02-10-2002
			DK 721777 T3	07-10-2002
			EP 0721777 A2	17-07-1996
			ES 2181845 T3	01-03-2003
			FI 972922 A	09-07-1997
			HU 9801283 A2	28-06-1999
			JP 10512262 T	24-11-1998
			NO 973170 A	02-09-1997
			NZ 301500 A	28-07-2000
			PL 321273 A1	24-11-1997
			PT 721777 T	29-11-2002
			RU 2163802 C2	10-03-2001
			TR 9700627 T1	21-01-1998
			WO 9621430 A1	18-07-1996
WO 02053140	A	11-07-2002	WO 02053140 A2	11-07-2002
			US 2002156067 A1	24-10-2002
WO 02053104	A	11-07-2002	WO 02053104 A2	11-07-2002
			US 2002161002 A1	31-10-2002
WO 0127068	A	19-04-2001	AU 7307000 A	23-04-2001
			BG 106603 A	29-12-2002
			BR 0014733 A	11-06-2002
			CN 1378527 T	06-11-2002
			CZ 20021180 A3	13-11-2002
			EP 1220831 A1	10-07-2002
			WO 0127068 A1	19-04-2001
			NO 20021659 A	08-04-2002
			TR 200201004 T2	21-11-2002
			US 6410736 B1	25-06-2002
EP 0534756	A	31-03-1993	US 5281624 A	25-01-1994

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/36132

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0534756	A	AT 124934 T	15-07-1995
		AU 655075 B2	01-12-1994
		AU 2537092 A	01-04-1993
		BR 1100384 A3	31-08-1999
		BR 9203743 A	20-04-1993
		CA 2079161 A1	28-03-1993
		CA 2079162 A1	28-03-1993
		CN 1071416 A ,B	28-04-1993
		CZ 9202908 A3	14-04-1993
		DE 69203424 D1	17-08-1995
		DE 69203424 T2	11-01-1996
		DK 537915 T3	27-11-1995
		EP 0534756 A2	31-03-1993
		EP 0537915 A1	21-04-1993
		EP 0591584 A1	13-04-1994
		FI 924305 A	28-03-1993
		HU 65375 A2	02-05-1994
		IL 103277 A	20-11-1997
		JP 3256291 B2	12-02-2002
		JP 5201961 A	10-08-1993
		JP 5238996 A	17-09-1993
		KR 221180 B1	15-09-1999
		MX 9205433 A1	01-03-1993
		NO 923710 A ,B,	29-03-1993
		NZ 244460 A	26-10-1994
		SK 290892 A3	14-02-2000
		RU 2057120 C1	27-03-1996
		ZA 9207240 A	22-03-1994